Quiz for Ostomy Medication Administration

NAM	E		DATE	SCHOOL	
1.	When resistance is present when administering an ostomy medication, what should you do?				
	a.	Call 911, this is a medical e	mergency.		
	b.	This is a common problem;		past the resistance.	
	c.	Check to make sure the clar	np is open. If clamp is open a	nd you still feel resistance, call school nurse and/or	
		parent to intervene as soon a	as possible.		
	d.	Eliminate the feeding and w	rite a note to the parent expla	ining your difficulty.	
2.	If air bubbles are present when drawing up the medication into the syringe, what is your next step?				
	a.				
	b.		in the syringe and start over		
	c.				
	d.	inadequate amount, drawn up more medication into the syringe to the prescribed dosage. Add more medication to the syringe to account for the air bubble.			
3.	It is important to make sure that the student's head is raised at least 30 degrees when administering medication through thei gastrostomy tube?				
		True			
	b.	False			
4.	What is the best sequence of events when administering an ostomy medication?				
	a.	do, recheck the first five rig	hts, administer the medication	pour the medication, tell student what you are going n, remove gloves and wash hands, record on	
	1	medication administration r			
	b.		re going to do, record on medinister the medication, remo	cation administration record, check the first five	
	c.			and apply gloves, recheck the first five rights, tell	
	C.			dministration record, administer the medication,	
		remove gloves and wash ha		,	
	d.			ration record, wash hands and apply gloves, tell	
5.	Which of th			on, remove gloves and wash hands. or medication administration?	
		•			
	a.	Right equipment			
	b.	Right medication			
	c. d.	Right student Right time			
	u.	right time			

Number correct: _____ of 5 answers