

Quiz for General Principles and Oral Medication Administration

Please circle the correct answer and give the completed test to the school nurse for scoring and to schedule hands-on training.

NAME _____ DATE _____ SCHOOL _____

1. Which of the following medication guidelines or rights are not appropriate for medication administration?
 - a. Right student
 - b. Right dosage
 - c. Right place
 - d. Right time

2. Which of the following is considered a medication error?
 - a. Student forgets to come to the health office for daily medication.
 - b. Medication is not available at school due to inability of the parents to bring in additional medication.
 - c. Student refuses to take the daily medication.
 - d. Student vomits the medication 15 minutes after administration of the medication.

3. When administering liquid medications which of the following are true?
 - a. All medication should be shaken before pouring in a measuring cup.
 - b. Only elixir medication should be shaken prior to pouring in a measuring cup.
 - c. When pouring liquid medication into a measuring cup use the sides of the meniscus to measure the correct amount of medication.
 - d. Only suspension medications should be shaken using the bottom of the meniscus in the measuring cup to measure the correct amount of medication.

4. What documentation is necessary prior to the administration of prescription medications?
 - a. Medical provider authorization and parent consent
 - b. Properly labeled prescription bottle
 - c. Parent consent and properly labeled prescription bottle
 - d. Medical provider and properly labeled prescription bottle

5. School personnel should check the first five rights or guidelines for medication administration each of the times below except?
 - a. Before pouring the medication into a medication cup
 - b. Upon arrival of the medication.
 - c. Prior to administration of the medication to the student
 - d. After administration of the medication to the student.

Number correct: _____ of 5 answers